

PRINCIPAL ARC ANALYSIS ON DIRECT PRODUCT MANIFOLDS

BY SUNGKYU JUNG, MARK FOSKEY AND J. S. MARRON

*University of North Carolina, University of North Carolina and
 University of North Carolina*

We propose a new approach to analyze data that naturally lie on manifolds. We focus on a special class of manifolds, called direct product manifolds, whose intrinsic dimension could be very high. Our method finds a low-dimensional representation of the manifold that can be used to find and visualize the principal modes of variation of the data, as Principal Component Analysis (PCA) does in linear spaces. The proposed method improves upon earlier manifold extensions of PCA by more concisely capturing important nonlinear modes. For the special case of data on a sphere, variation following nongeodesic arcs is captured in a single mode, compared to the two modes needed by previous methods. Several computational and statistical challenges are resolved. The development on spheres forms the basis of principal arc analysis on more complicated manifolds. The benefits of the method are illustrated by a data example using medial representations in image analysis.

1. Introduction. Principal Component Analysis (PCA) has been frequently used as a method of dimension reduction and data visualization for high-dimensional data. For data that naturally lie in a curved manifold, application of PCA is not straightforward since the sample space is not linear. Nevertheless, the need for PCA-like methods is growing as more manifold data sets are encountered and as the dimensions of the manifolds increase.

In this article we introduce a new approach for an extension of PCA on a special class of manifold data. We focus on direct products of simple manifolds, in particular, of the unit circle S^1 , the unit sphere S^2 , \mathbb{R}_+ and \mathbb{R}^p . We will refer to these as *direct product manifolds*, for convenience. Many types of statistical sample spaces are special cases of the direct product manifold. A widely known example is the sample space for directional data [Fisher (1993), Fisher, Lewis and Embleton (1993) and Mardia and Jupp (2000)] and their direct products. Applications include analysis of wind directions, orientations of cracks, magnetic field directions and directions from the earth to celestial objects. For example, when we consider multiple 3-D directions simultaneously, the sample space is $S^2 \otimes \cdots \otimes S^2$, which is a direct product manifold. Another example is the medial representation of shapes [m-reps, Siddiqi and Pizer (2008)], which is somewhat less known to the statistical community but provides a powerful parametrization of 3-D shapes

Received December 2009; revised May 2010.

Key words and phrases. Principal Component Analysis, nonlinear dimension reduction, manifold, folded Normal distribution, directional data, image analysis, medial representation.

1 of human organs and has been extensively studied in the image analysis field. 1
2 The space of m-reps is usually a high-dimensional direct product manifold. Some 2
3 background and necessary definitions on direct product manifolds can be found in 3
4 [Appendix](#). 4

5 Our approach to a manifold version of PCA builds upon earlier work, espe- 5
6 cially the principal geodesic analysis proposed by [Fletcher et al. \(2004\)](#) and the 6
7 geodesic PCA proposed by [Huckemann and Ziezold \(2006\)](#) and [Huckemann, Hotz 7
8 and Munk \(2010\)](#). A detailed catalogue of current methodologies can be found in 8
9 [Huckemann, Hotz and Munk \(2010\)](#). An important approach among these is to 9
10 approximate the manifold by a linear space. [Fletcher et al. \(2004\)](#) take the tangent 10
11 space of the manifold at the geodesic mean as the linear space, and work with 11
12 appropriate mappings between the manifold and the tangent space. This results in 12
13 finding the best fitting geodesics among those passing through the geodesic mean. 13
14 This was improved in an important way by [Huckemann, Hotz and Munk](#), who 14
15 found the best fit over the set of all geodesics. [Huckemann, Hotz and Munk](#) went 15
16 on to propose a new notion of center point, the *PCmean*, which is an intersection 16
17 of the first two principal geodesics. This approach gives significant advantages, es- 17
18 pecially when the curvature of the manifold makes the geodesic mean inadequate, 18
19 an example of which is depicted in [Figure 2b](#). 19

20 Our method inherits advantages of these methods and improves further by ef- 20
21 fectively capturing more complex nongeodesic modes of variation. Note that the 21
22 curvature of direct product manifolds is mainly due to the spherical part, which 22
23 motivates careful investigation of S^2 -valued variables. We point out that (small) 23
24 circles in S^2 , including geodesics, can be used to capture the nongeodesic varia- 24
25 tion. We introduce the *principal circles* and *principal circle mean*, analogous to, 25
26 yet more flexible than, the geodesic principal component and PCmean of [Hucke- 26
27 mann, Hotz and Munk](#). These become *principal arcs* when the manifold is indeed 27
28 S^2 . For more complex direct product manifolds, we suggest transforming the data 28
29 points in S^2 into a linear space by a special mapping utilizing the principal cir- 29
30 cles. For the other components of the manifold, the tangent space mappings can 30
31 be used to map the data into a linear space as done in [Fletcher et al. \(2004\)](#). Once 31
32 manifold-valued data are mapped onto the linear space, then the classical linear 32
33 PCA can be applied to find principal components in the transformed linear space. 33
34 The estimated principal components in the linear space can be back-transformed 34
35 to the manifold, which leads to principal arcs. 35

36 We illustrate the potential of our method by an example of m-rep data in [Fig- 36
37 ure 1](#). Here, m-reps with 15 sample points called atoms model the prostate gland 37
38 (an organ in the male reproductive system) and come from the simulator developed 38
39 and analyzed in [Jeong et al. \(2008\)](#). [Figure 1](#) shows that the S^2 components of the 39
40 data tend to be distributed along small circles, which frequently are not geodesics. 40
41 We emphasize the curvature of variation along each sphere by fitting a great circle 41
42 and a small circle (by the method discussed in [Section 2](#)). Our method is adapted 42
43 43

PRINCIPAL ARC ANALYSIS

3




FIG. 1. S^2 -valued samples ($n = 60$) of the prostate *m-reps* with 15 medial atoms. One sample in this figure is represented by a 30-tuple of 3-D directions (two directions at each atom), which lies in the manifold $\otimes_{i=1}^{15} (S^2 \otimes S^2)$. Small and great circles are fitted and plotted in the rightmost atoms to emphasize the sample variation along small circles.

to capture this nonlinear (nongeodesic) variation of the data. A potential application of our method is to improve accuracy of segmentation of objects from CT images. Detailed description of the data and results of our analysis can be found in Section 5.

Note that the previous approaches [Fletcher et al. (2004), Huckemann and Ziezold (2006)] are defined for general manifolds, while our method focuses on these particular direct product manifolds. Although the method is not applicable for general manifolds, it is useful for this common class of manifolds that is often found in applications. Our results inform our belief that focusing on specific types of manifolds allow more precise and informative statistical modeling than methods that attempt to be fully universal. This happens through using special properties (e.g., presence of small circles) that are not available for all other manifolds.

The rest of the article is organized as follows. We begin by introducing a *circle class* on S^2 as an alternative to the set of geodesics. Section 2 discusses principal circles in S^2 , which will be the basis of the special transformation. The first principal circle is defined by the least-squares circle, minimizing the sum of squared residuals. In Section 3 we introduce a data-driven method to decide whether the least-squares circle is appropriate. A recipe for principal arc analysis on direct product manifolds is proposed in Section 4 with discussion on the transformations. A detailed introduction of the space of *m-reps* and the results from applying the proposed method follow. A novel computational algorithm for the least-squares circles is presented in Section 6. In the Appendix we provide some necessary background for treating direct product manifolds as sample spaces, including the notion of geodesic mean, tangent space, exponential map and log map.

2. Circle class for nongeodesic variation on S^2 . Consider a set of points in \mathbb{R}^2 . Numerous methods for understanding population properties of a data set in linear space have been proposed and successfully applied, which include rigid methods, such as linear regression and principal components, and very flexible methods, such as scatterplot smoothing and principal curves [Hastie and Stuetzle (1989)]. In this paper we make use of a parametric class of circles, including small and great circles, which allows much more flexibility than either methods of Fletcher et al. (2004) or Huckemann, Hotz and Munk (2010), but less flexibility than a principal curve approach. Although this idea was motivated by examples such as those in Figure 1, there are more advantages gained from using the class of circles:

- (i) The circle class includes the simple geodesic case.
- (ii) Each circle can be parameterized, which leads to an easy interpretation.
- (iii) There is an orthogonal complement of each circle, which gives two important advantages:
 - (a) Two orthogonal circles can be used as a basis of a further extension to principal arc analysis.
 - (b) Building a sensible notion of principal components on S^2 alone is easily done by utilizing the circles.

The idea (iii)(b) will be discussed in detail after introducing a method of circle fitting. Some notations follow: S^2 can be thought of as the unit sphere in \mathbb{R}^3 , so that a unit vector $\mathbf{x} \in \mathbb{R}^3$ is a member of S^2 . The geodesic distance between $\mathbf{x}, \mathbf{y} \in S^2$, denoted by $\rho(\mathbf{x}, \mathbf{y})$, is defined as the shortest distance between \mathbf{x}, \mathbf{y} along the sphere, which is the same as the angle formed by the two vectors. Thus, $\rho(\mathbf{x}, \mathbf{y}) = \cos^{-1}(\mathbf{x}'\mathbf{y})$. A circle on S^2 is conveniently parameterized by center $\mathbf{c} \in S^2$ and geodesic radius r , and denoted by $\delta(\mathbf{c}, r) = \{\mathbf{x} \in S^2 | \rho(\mathbf{c}, \mathbf{x}) = r\}$. It is a geodesic when $r = \pi/2$. Otherwise it is a small circle.

A circle that best fits the points $\mathbf{x}_1, \dots, \mathbf{x}_n \in S^2$ is found by minimizing the sum of squared residuals. The residual of \mathbf{x}_i is defined as the signed geodesic distance from \mathbf{x}_i to the circle $\delta(\mathbf{c}, r)$. Then the least-squares circle is obtained by

$$(1) \quad \min_{\mathbf{c}, r} \sum_{i=1}^n (\rho(\mathbf{x}_i, \mathbf{c}) - r)^2 \quad \text{subject to } \mathbf{c} \in S^2, r \in (0, \pi).$$

Note that there are always multiple solutions of (1). In particular, whenever (\mathbf{c}, r) is a solution, $(-\mathbf{c}, \pi - r)$ also solves the problem as $\delta(\mathbf{c}, r) = \delta(-\mathbf{c}, \pi - r)$. This ambiguity does not affect any essential result in this paper. Our convention is to use the circle with smaller geodesic radius.

The optimization task (1) is a constrained nonlinear least squares problem. We propose an algorithm to solve the problem that features a simplified optimization task and approximation of S^2 by tangent planes. The algorithm works in a doubly iterative fashion, which has been shown by experience to be stable and fast. Section 6 contains a detailed illustration of the algorithm.

1 Analogous to principal geodesics in S^2 , we can define *principal circles* in S^2 by 1
 2 utilizing the least-squares circle. The principal circles are two orthogonal circles 2
 3 in S^2 that best fit the data. We require the first principal circle to minimize the 3
 4 variance of the residuals, so it is the least-squares circle (1). The second principal 4
 5 circle is a geodesic which passes through the center of the first circle and thus is 5
 6 orthogonal at the points of intersection. Moreover, the second principal circle is 6
 7 chosen so that one intersection point is the intrinsic mean [defined in (2) later] of 7
 8 the projections of the data onto the first principal circle. 8

9 Based on a belief that the intrinsic (or extrinsic) mean defined on a curved 9
 10 manifold may not be a useful notion of the center point of the data [see, e.g., 10
 11 [Huckemann, Hotz and Munk](#) and Figure 2b], the principal circles do not use the 11
 12 pre-determined means. To develop a better notion of center point, we locate the 12
 13 best 0-dimensional representation of the data in a data-driven manner. Inspired by 13
 14 the PCmean idea of [Huckemann, Hotz and Munk](#), given the first principal circle 14
 15 δ_1 , the *principal circle mean* $\mathbf{u} \in \delta_1$ is defined (in an intrinsic way) as 15

$$16 \quad (2) \quad \mathbf{u} = \operatorname{argmin}_{\mathbf{u} \in \delta_1} \sum_{i=1}^n \rho^2(\mathbf{u}, P_{\delta_1} \mathbf{x}_i), \quad 16$$

17 where $P_{\delta_1} \mathbf{x}$ is the projection of \mathbf{x} onto δ_1 , that is, the point on δ_1 of the shortest 17
 18 geodesic distance to \mathbf{x} . Then 18

$$19 \quad (3) \quad P_{\delta_1(\mathbf{c}, r)} \mathbf{x} = \frac{\mathbf{x} \sin(r) + \mathbf{c} \sin(\rho(\mathbf{x}, \mathbf{c}) - r)}{\sin(\rho(\mathbf{x}, \mathbf{c}))}, \quad 19$$

20 as in equation (3.3) of [Mardia and Gadsden \(1977\)](#). We assume that \mathbf{c} is the north 20
 21 pole \mathbf{e}_3 , without losing generality since otherwise the sphere can be rotated. Then 21

$$22 \quad (4) \quad \rho(\mathbf{u}, P_{\delta_1} \mathbf{x}) = \sin(r) \rho_{S^1} \left(\frac{(u_1, u_2)}{\sqrt{1 - u_3^2}}, \frac{(x_1, x_2)}{\sqrt{1 - x_3^2}} \right), \quad 22$$

23 where $\mathbf{u} = (u_1, u_2, u_3)'$, $\mathbf{x} = (x_1, x_2, x_3)'$ and ρ_{S^1} is the geodesic (angular) distance 23
 24 function on S^1 . The optimization problem (2) is equivalent to finding the geodesic 24
 25 mean in S^1 . See equation (17) in the [Appendix](#) for computation of the geodesic 25
 26 mean in S^1 . 26

27 The second principal circle δ_2 is then the geodesic passing through the principal 27
 28 circle mean \mathbf{u} and the center \mathbf{c} of δ_1 . Denote $\bar{\delta} \equiv \bar{\delta}(\mathbf{x}_1, \dots, \mathbf{x}_n)$ as a combined 28
 29 representation of (δ_1, \mathbf{u}) or, equivalently, (δ_1, δ_2) . 29

30 As a special case, we can force the principal circles to be great circles. The 30
 31 best fitting geodesic is obtained as a solution of the problem (1) with $r = \pi/2$ 31
 32 and becomes the first principal circle. The optimization algorithm for this case 32
 33 is slightly modified from the original algorithm for the varying r case by simply 33
 34 setting $r = \pi/2$. The principal circle mean \mathbf{u} and the δ_2 for this case are defined 34
 35 in the same way as in the small circle case. Note that the principal circles with 35
 36 $r = \pi/2$ are essentially the same as the method of [Huckemann and Ziezold \(2006\)](#). 36
 37 37
 38 38
 39 39
 40 40
 41 41
 42 42
 43 43




FIG. 2. Toy examples on S^2 with $n = 30$ points showing the first principal circle (red) as a small circle and the first geodesic principal component (dotted green) by Huckemann, and the first principal geodesic (black) by Fletcher. Also plotted are the geodesic mean, PCmean and principal circle mean of the data as black, green and red diamonds, respectively. (a) The three methods give similar satisfactory answers when the data are stretched along a geodesic. (b) When the data are stretched along a great circle, covering almost all of it, the principal geodesic (black circle) and geodesic mean (black diamond) fail to find a reasonable representation of the data, while the principal circle and Huckemann's geodesic give sensible answers. (c) Only the principal circle fits well when the data are not along a geodesic. (d) For a small cluster without principal modes of variation, the principal circle gets too small. See Section 3 for discussion of this phenomenon.

Figure 2 illustrates the advantages of using the circle class to efficiently summarize variation. On four different sets of toy data, the first principal circle δ_1 is plotted with principal circle mean \mathbf{u} . The first principal geodesics from the methods of Fletcher and Huckemann are also plotted with their corresponding mean. Figure 2a illustrates the case where the data were indeed stretched along a geodesic. The solutions from the three methods are similar to one another. The advantage of Huckemann's method over Fletcher's can be found in Figure 2b. The geodesic mean is found far from the data, which leads to poor performance of the principal geodesic analysis, because it considers only great circles passing through the geo-

1 desic mean. Meanwhile, the principal circle and Huckemann’s method, which do 1
 2 not utilize the geodesic mean, work well. The case where geodesic mean and any 2
 3 geodesic do not fit the data well is illustrated in Figure 2c, which is analogous to 3
 4 the Euclidean case, where a nonlinear fitting may do a better job of capturing the 4
 5 variation than PCA. To this data set, the principal circle fits best, and our defini- 5
 6 tion of mean is more sensible than the geodesic mean and the PCmean. The points 6
 7 in Figure 2d are generated from the von Mises–Fisher distribution with $\kappa = 10$, 7
 8 thus having no principal mode of variation. In this case the first principal circle 8
 9 δ_1 follows a contour of the apparent density of the points. We shall discuss this 9
 10 phenomenon in detail in the following section. 10

11 Fitting a (small) circle to data on a sphere has been investigated for some time, 11
 12 especially in statistical applications in geology. Those approaches can be distin- 12
 13 guished in three different ways, where our choice fits into the first category: 13
 14

- 15 (1) Least-squares of intrinsic residuals: Gray, Geiser and Geiser (1980) formulated 15
 16 the same problem as in (1), finding a circle that minimizes sum of squared 16
 17 residuals, where residuals are defined in a geodesic sense. 17
- 18 (2) Least-squares of extrinsic residuals: A different measure of residual was cho- 18
 19 sen by Mardia and Gadsden (1977) and Rivest (1999), where the residual of 19
 20 \mathbf{x} from $\delta(c, r)$ is defined by the shortest Euclidean distance between \mathbf{x} and 20
 21 $\delta(\mathbf{c}, r)$. Their objective is to find 21

$$22 \quad \operatorname{argmin}_{\delta} \sum_{i=1}^n \|\mathbf{x}_i - P_{\delta} \mathbf{x}_i\|^2 = \operatorname{argmin}_{\delta} \sum_{i=1}^n -\mathbf{x}_i' P_{\delta} \mathbf{x}_i = \operatorname{argmin}_{\delta} \sum_{i=1}^n -\cos(\xi_i), \quad 22$$

23 where ξ_i denotes the intrinsic residual. This type of approach can be numeri- 25
 26 cally close to the intrinsic method as $\cos(\xi_i) = 1 - \xi_i^2/2 + O(\xi_i^4)$. 26

- 27 (3) Distributional approach: Mardia and Gadsden (1977) and Bingham and Mar- 27
 28 dia (1978) proposed appropriate distributions to model S^2 -valued data that 28
 29 cluster near a small circle. These models essentially depend on the quantity 29
 30 $\cos(\xi)$, which is easily interpreted in the extrinsic sense but not in the intrinsic 30
 31 sense. 31
 32

33 **REMARK.** The principal circle and principal circle mean always exist. This is 33
 34 because the objective function (1) is a continuous function of \mathbf{c} , with the compact 34
 35 domain S^2 . The minimizer r has a closed-form solution (see Section 6). A similar 35
 36 argument can be made for the existence of \mathbf{u} . On the other hand, the uniqueness of 36
 37 the solution is not guaranteed. We conjecture that if the manifold is approximately 37
 38 linear or, equivalently, the data set is well approximated by a linear space, then the 38
 39 principal circle will be unique. However, this does not lead to the uniqueness of \mathbf{u} , 39
 40 whose sufficient condition is that the projected data on δ_1 is strictly contained in a 40
 41 half-circle [Karcher (1977)]. Note that a sufficient condition for the uniqueness of 41
 42 the principal circle is not clear even in the Euclidean case [Chernov (2010)]. 42
 43

3. Suppressing small least-squares circles. When the first principal circle δ_1 has a small radius, sometimes it is observed that δ_1 does not fit the data in a manner that gives useful decomposition, as shown in Figure 2d. This phenomenon has been also observed for the related principal curve fitting method of [Hastie and Stuetzle \(1989\)](#). We view this as unwanted overfitting, which is indeed a side effect caused by using the full class of circles with free radius parameter instead a class of great circles. In this section a data-driven method to flag this overfitting is discussed. In essence, the fitted small circle is replaced by the best fitting geodesics when the data do not cluster along the circle but instead tend to cluster near the center of the circle.

We first formulate the problem and solution in \mathbb{R}^2 . This is for the sake of clear presentation and also because the result on \mathbb{R}^2 can be easily extended to S^2 using a tangent plane approximation.

Let $f_{\mathbf{X}}$ be a spherically symmetric density function of a continuous distribution defined on \mathbb{R}^2 . Whether the density is high along some circle is of interest. By the symmetry assumption, density height along a circle can be found by inspecting a section of $f_{\mathbf{X}}$ along a ray from the origin (the point of symmetry). A section of $f_{\mathbf{X}}$ coincides with the conditional density $f_{X_1|X_2}(x_1|x_2=0) = \kappa^{-1} f_{\mathbf{X}}(x_1, 0)$. A random variable corresponding to the p.d.f. $f_{X_1|X_2=0}$ is not directly observable. Instead, the radial distance $R = \|\mathbf{X}\|$ from the origin can be observed. For the polar coordinates (R, Θ) such that $\mathbf{X} = (X_1, X_2) = (R \cos \Theta, R \sin \Theta)$, the marginal p.d.f. of R is $f_R(r) = 2\pi r f_{\mathbf{X}}(r, 0)$ as $f_{\mathbf{X}}$ is spherically symmetric. A section of $f_{\mathbf{X}}$ is related to the observable density f_R as $f_R(r) \propto r f_{X_1|X_2=0}(r)$, for $r \geq 0$. This relation is called the length-biased sampling problem [[Cox \(1969\)](#)]. The relation can be understood intuitively by observing that a value r of R can be observed at any point on a circle of radius r , circumference of which is proportional to r . Thus, sampling of R from the density $f_{X_1|X_2=0}$ is proportional to its size.

The problem of suppressing a small circle can be paraphrased as “how to determine whether a nonzero point is a mode of the function $f_{X_1|X_2=0}$, when observing only a length-biased sample.”

The spectrum from the circle-clustered case (mode at a nonzero point) to the center-clustered case (mode at origin) can be modeled as

$$(5) \quad \text{data} = \text{signal} + \text{error},$$

where the signal is along a circle with radius μ , and the error accounts for the perpendicular deviation from the circle (see Figure 3). Then, in polar coordinates (R, Θ) , Θ is uniformly distributed on $(0, 2\pi]$ and R is a positive random variable with mean μ . First assume that R follows a truncated Normal distribution with standard deviation σ , with the marginal p.d.f. proportional to

$$(6) \quad f_R(r) \propto \phi\left(\frac{r - \mu}{\sigma}\right) \quad \text{for } r \geq 0,$$




FIG. 3. Illustration of the conceptual model (5) on \mathbb{R}^2 , which can also be understood as a local approximation of S^2 . The signal is along the circle centered at c and radius μ . The error is perpendicular to the signal. When the deviation σ is large, it is possible that the amount of error is even greater than the radius μ . This is incorporated in the wrapping approach (7).

where ϕ is the standard Normal density function. The conditional density $f_{X_1|X_2=0}$ is then

$$f_{X_1|X_2=0}(r) \propto \frac{1}{r} f_R(r) \propto \frac{1}{r\sigma} \exp\left(-\frac{(r-\mu)^2}{\sigma^2}\right) \quad \text{for } r > 0.$$

Nonzero local extrema of $f_{X_1|X_2=0}$ can be characterized as a function of (μ, σ) in terms of $r_+, r_- = \{\mu \pm \sqrt{(\mu-2\sigma)(\mu+2\sigma)}\}/2$ as follows:

- When $\mu > 2\sigma$, $f_{X_1|X_2=0}$ has a local maximum at r_+ , minimum at r_- .
- When $\mu = 2\sigma$, $r_+ = r_- = \frac{\mu}{2}$.
- When $\mu < 2\sigma$, $f_{X_1|X_2=0}$ is strictly decreasing, for $r \geq 0$.

Therefore, whenever the ratio $\mu/\sigma > 2$, $f_{X_1|X_2=0}$ has a mode at r_+ .

This idea can be applied for circles in S^2 with some modification, shown next. We point out that the model (5) is useful for understanding the small circle fitting: signal as a circle with radius μ , and error as the deviation along geodesics perpendicular to the circle. Moreover, a spherically symmetric distribution centered at \mathbf{c} on S^2 can be mapped to a spherically symmetric distribution on the tangent space at \mathbf{c} , preserving the radial distances by the log map (defined in the Appendix). A modification needs to be made on the truncated density f_R . It is more natural to let the error be so large that the deviation from the great circle is greater than μ . Then the observed value may be found near the opposite side of the true signal, which is illustrated in Figure 3 as the large deviation case. To incorporate this case, we consider a wrapping approach. The distribution of errors (on the real line) is wrapped around the sphere along a great circle through \mathbf{c} , and the marginal p.d.f. f_R in (6) is modified to

$$(7) \quad f_R^w(r) \propto \sum_{k=0}^{\infty} \left[\phi\left(\frac{r+2\pi k-\mu}{\sigma}\right) + \phi\left(\frac{r-2\pi k+\mu}{\sigma}\right) \right] \quad \text{for } r \in [0, \pi].$$




FIG. 4. (Left) Graph of $f_{X_1|X_2=0}^w(r)$ for $\mu/\sigma = 1, 2, 3, 4$. The density is high at a nonzero point when $\mu/\sigma > 2.0534$. (Center, right) Spherically symmetric distributions f corresponding to $\mu/\sigma = 2, 3$. The ratio $\mu/\sigma > 2$ roughly leads to a high density along a circle.

The corresponding conditional p.d.f., $f_{X_1|X_2=0}^w$, is similar to $f_{X_1|X_2=0}$ and a numerical calculation shows that $f_{X_1|X_2=0}^w$ has a mode at some nonzero point whenever $\mu/\sigma > 2.0534$, for $\mu < \pi/2$. In other words, we use the small circle when μ/σ is large. Note that in what follows we only consider the first term ($k = 0$) of (7) since other terms are negligible in most situations. We have plotted $f_{X_1|X_2=0}^w$ for some selected values of μ and σ in Figure 4.

With a data set on S^2 , we need to estimate μ and σ , or the ratio μ/σ . Let $\mathbf{x}_1, \dots, \mathbf{x}_n \in S^2$ and let $\hat{\mathbf{c}}$ be the samples and the center of the fitted circle, respectively. Denote ξ_i for the errors of the model (5) such that $\xi_i \sim N(0, \sigma^2)$. Then $r_i \equiv \rho(\mathbf{x}_i, \hat{\mathbf{c}}) = |\mu + \xi_i|$, which has the *folded normal* distribution [Leone, Nelson and Nottingham (1961)]. Estimation of μ and σ based on unsigned r_i is not straightforward. We present two different approaches to this problem.

Robust approach. The observations r_1, \dots, r_n can be thought of as a set of positive numbers contaminated by the folded negative numbers. Therefore, the left half (near zero) of the data are more contaminated than the right half. We only use the right half of the data, which are less contaminated than the other half. We propose to estimate μ and σ by

$$(8) \quad \hat{\mu} = \text{med}(r_1^n), \quad \hat{\sigma} = (Q_3(r_1^n) - \text{med}(r_1^n))/Q_3(\Phi),$$

where $Q_3(\Phi)$ is the third quantile of the standard normal distribution. The ratio can be estimated by $\hat{\mu}/\hat{\sigma}$.

Likelihood approach via EM algorithm. The problem may also be solved by a likelihood approach. Early solutions can be found in Leone, Nelson and Nottingham (1961), Elandt (1961) and Johnson (1962), in which the MLEs were given by numerically solving nonlinear equations based on the sample moments. As those methods were very complicated, we present a simpler approach based on the EM algorithm. Consider unobserved binary variables s_i with values -1 and $+1$ so that $s_i r_i \sim N(\mu, \sigma^2)$. The idea of the EM algorithm is that if we have observed s_i , then

1 the maximum likelihood estimator of $\vartheta = (\mu, \sigma^2)$ would be easily obtained. The 1
 2 EM algorithm is an iterative algorithm consisting of two steps. Suppose that the 2
 3 k th iteration produced an estimate $\hat{\vartheta}_k$ of ϑ . The E-step is to impute s_i based on r_i 3
 4 and $\hat{\vartheta}_k$ by forming a conditional expectation of log-likelihood for ϑ , 4

$$\begin{aligned}
 5 \quad Q(\vartheta) &= E \left[\log \prod_{i=1}^n f(r_i, s_i | \vartheta) \middle| r_i, \hat{\vartheta}_k \right] \\
 6 \\
 7 \\
 8 \\
 9 \quad &= \sum_{i=1}^n [\log f(r_i | s_i = +1, \vartheta) P(s_i = +1 | r_i, \hat{\vartheta}_k) \\
 10 \\
 11 \quad &\quad + \log f(r_i | s_i = -1, \vartheta) P(s_i = -1 | r_i, \hat{\vartheta}_k)] \\
 12 \\
 13 \quad &= \sum_{i=1}^n [\log \phi(r_i | \vartheta) p_{i(k)} + \log \phi(-r_i | \vartheta) (1 - p_{i(k)})], \\
 14 \\
 15
 \end{aligned}$$

16 where f is understood as an appropriate density function, and $p_{i(k)}$ is easily com- 16
 17 puted as 17

$$18 \quad p_{i(k)} = P(s_i = +1 | r_i, \hat{\vartheta}_k) = \frac{\phi(r_i | \hat{\vartheta}_k)}{\phi(r_i | \hat{\vartheta}_k) + \phi(-r_i | \hat{\vartheta}_k)}. \quad 18$$

19 The M-step is to maximize $Q(\vartheta)$ whose solution becomes the next estimator $\hat{\vartheta}_{k+1}$. 19
 20 Now the $(k + 1)$ th estimates are calculated by a simple differentiation and given 20
 21 by 21

$$22 \quad \hat{\mu}_{k+1} = \frac{1}{n} \sum_{i=1}^n (2p_{i(k)} - 1)r_i, \quad \hat{\sigma}_{k+1}^2 = \frac{1}{n} \sum_{i=1}^n (r_i^2 - \hat{\mu}_{k+1}^2). \quad 22$$

23 With the sample mean and variance of r_1, \dots, r_n as an initial estimator $\hat{\vartheta}_0$, the 23
 24 algorithm iterates E-steps and M-steps until the iteration changes the estimates 24
 25 less than a predefined criteria (e.g., 10^{-10}). μ/σ is estimated by the ratio of the 25
 26 solutions. 26

27 *Comparison.* Performance of these estimators are now examined by a simula- 27
 28 tion study. Normal random samples are generated with ratios μ/σ being 0, 1, 2 28
 29 or 3, representing the transition from the center-clustered to circle-clustered case. 29
 30 For each ratio, $n = 50$ samples are generated, from which $\hat{\mu}/\hat{\sigma}$ is estimated. These 30
 31 steps are repeated 1000 times to obtain the sampling variation of the estimates. 31
 32 We also study the $n = 1000$ case in order to investigate the consistency of the 32
 33 estimators. The results are summarized in Figure 5 and Table 1. 33

34 The distribution of estimators are shown for $n = 50, 1000$ in Figure 5 and the 34
 35 proportion of estimators greater than 2 is summarized in Table 1. When $n = 1000$, 35
 36 both estimators are good in terms of the proportion of correct answers. In the fol- 36
 37 lowing, the proportions of correct answers are corresponding to $n = 50$ case. The 37
 38 38
 39 39
 40 40
 41 41
 42 42
 43 43




FIG. 5. Simulation results of the proposed estimators for the ratio μ/σ . Different ratios represent different underlying distributions. For example, estimators in the top left are based on random samples from a folded Normal distribution with mean $\mu = 3$, standard deviation $\sigma = 1$. Curves are smooth histograms of estimates from 1000 repetitions. The thick black curve represents the distribution of the robust estimator from $n = 50$ samples. Likewise, the thick red curve is for the MLE with $n = 50$, the dotted black curve is for the robust estimator with $n = 1000$, and the dotted red curve is for the MLE with $n = 1000$. The smaller sample size represents a usual data analytic situation, while the $n = 1000$ case shows an asymptotic situation.

top left panel in Figure 5 illustrates the circle-centered case with ratio 3. The estimated ratios from the robust approach give correct solutions (greater than 2) 95% of the time (98.5% for likelihood approach). For the borderline case (ratio 2, top right), the small circle will be used about half the time. The center-clustered case is demonstrated with the true ratio 1, that also gives a reasonable answer (proportion of correct answers 95.3% and 94.8% for the robust and likelihood answers respectively). It can be observed that when the true ratio is zero, the robust estimates are far from 0 (the bottom right in Figure 5). However, this is expected to occur because the proportion of uncontaminated data is low when the ratio is too small. However, those ‘inaccurate’ estimates are around 1 and less than 2 most of the time, which leads to ‘correct’ answers. The likelihood approach looks somewhat better with more hits near zero, but an asymptotic study [Johnson (1962)]

PRINCIPAL ARC ANALYSIS

13

TABLE 1

Proportion of estimates greater than 2 from the data illustrated in Figure 5. For $\mu/\sigma = 3$, shown are proportions of correct answers from each estimator. For $\mu/\sigma = 1$ or 0, shown are proportions of incorrect answers

Method	$\mu/\sigma = 3$	$\mu/\sigma = 2$	$\mu/\sigma = 1$	$\mu/\sigma = 0$
MLE, $n = 50$	98.5	55.2	5.2	6.8
Robust, $n = 50$	95.0	50.5	4.7	1.4
MLE, $n = 1000$	100	51.9	0	0
Robust, $n = 1000$	100	50.5	0	0

showed that the variance of the maximum likelihood estimator converges to infinity when the ratio tends to zero, as glimpsed in the long right tail of the simulated distribution.

In summary, we recommend use of the robust estimators (8), which are computationally light, straightforward and stable for all cases.

In addition, we point out that Gray, Geiser and Geiser (1980) and Rivest (1999) proposed to use a goodness-of-fit statistic to test whether the small circle fit is better than a geodesic fit. Let r_g and r_c be the sums of squares of the residuals from great and small circle fits. They claimed that $V = (n - 3)(r_g - r_c)/r_c$ is approximately distributed as $F_{1, n-3}$ for a large n if the great circle was true. However, this test does not detect the case depicted in Figure 2d. The following numerical example shows the distinction between our approach and the goodness-of-fit approach.

EXAMPLE 1. Consider the sets of data depicted in Figure 2. The goodness-of-fit test gives p -values of 0.51, 0.11359, 0 and 0.0008 for (a)–(d), respectively. The estimated ratios μ/σ are 14.92, 16.89, 14.52 and 1.55. Note that for (d), when the least-squares circle is too small, our method suggests to use a geodesic fit over a small circle while the goodness-of-fit test gives significance of the small circle. The goodness-of-fit method is not adequate to suppress the overfitting small circle in a way we desire.

A referee pointed out that the transition of the principal circle between great circle and small circle is not continuous. Specifically, when the data set is perturbed so that the principal circle becomes too small, then the principal circle and principal circle mean are abruptly replaced by a great circle and geodesic mean. As an example, we have generated a toy data set spread along a circle with some radial perturbation. The perturbation is continuously inflated, so that with large inflation, the data are no longer circle-clustered. In Figure 6 the $\widehat{\mu/\sigma}$ changes smoothly, but once the estimate hits 2 (our criterion), there is a sharp transition between small and great circles. Sharp transitions do naturally occur in the statistics of manifold data.




FIG. 6. (Left) The estimate $\widehat{\mu/\sigma}$ decreases smoothly as the perturbation is inflated. (Center, right) Snapshots of the toy data on a sphere. A very small perturbation of the data set leads to a sharp transition between the small circle (center) and great circle (right).

For example, even the simple geodesic mean can exhibit a major discontinuous transition resulting from an arbitrarily small perturbation of the data. However, the discontinuity between small and great circles does seem more arbitrary and thus may be worth addressing. An interesting open problem is to develop a blended version of our two solutions, for values of $\widehat{\mu/\sigma}$ near 2, which could be done by fitting circles with radii that are smoothly blended between the small circle radius and $\pi/2$.

4. Principal arc analysis on direct product manifolds. The discussions of the principal circles in S^2 play an important role in defining the principal arcs for data in a direct product manifold $M = M_1 \otimes M_2 \otimes \dots \otimes M_d$, where each M_i is one of the simple manifolds S^1, S^2, \mathbb{R}_+ and \mathbb{R} . We emphasize again that the curvature of the direct product manifold M is mainly due to the spherical components.

Consider a data set $x_1, \dots, x_n \in M$, where $x_i \equiv (x_i^1, \dots, x_i^d)$ such that $x_i^j \in M_j$. Denote $d_0 \geq d$ for the intrinsic dimension of M . The geodesic mean \bar{x} of the data is defined component-wise for each simple manifold M_j . Similarly, the tangent plane at \bar{x} , $T_{\bar{x}}M$, is also defined marginally, that is, $T_{\bar{x}}M$ is a direct product of tangent spaces of the simple manifolds. This tangent space gives a way of applying Euclidean space-based statistical methods by mapping the data onto $T_{\bar{x}}M$. We can manipulate this approximation of the data component-wise. In particular, the marginal data on the S^2 components can be represented in a linear space by a transformation $h_{\bar{\delta}}$, depending on the principal circles, that differs from the tangent space approximation.

Since the principal circles $\bar{\delta}$ capture the nongeodesic directions of variation, we use the principal circles as axes, which can be thought of as flattening the quadratic form of variation. In principle, we require a mapping $h_{\bar{\delta}}: S^2 \rightarrow \mathbb{R}^2$ to have the following properties: For $\bar{\delta} = (\delta_1, \delta_2) = ((\delta_1(\mathbf{c}, r), \mathbf{u})$:

- \mathbf{u} is mapped to the origin,
- δ_1 is mapped onto the x -axis, and

- 1 • δ_2 is mapped onto the y-axis. 1

2 Two reasonable choices of the mapping $h_{\bar{\delta}}$ will be discussed in Section 4.1 2
3 in detail. 3

4 The mapping $h_{\bar{\delta}}$ and the tangent space projection together give a linear space 4
5 representation of the data where the Euclidean PCA is applicable. The line seg- 5
6 ments corresponding to the sample principal component direction of the trans- 6
7 formed data can be mapped back to M , and become the principal arcs. 7

8 A procedure for principal arc analysis is as follows: 8
9

- 10 (1) For each j such that M_j is S^2 , compute principal circles $\bar{\delta} = \bar{\delta}(x_1^j, \dots, x_2^j)$ 10
11 and the ratio $\widehat{\mu/\sigma}$. If the ratio is greater than the predetermined value $\varepsilon = 2$, 11
12 then $\bar{\delta}$ is adjusted to be great circles as explained in Section 2. 12
13 (2) Let $\mathbf{h}: M \rightarrow \mathbb{R}^{d_0}$ be a transformation $\mathbf{h}(x) = (h_1(x^1), \dots, h_d(x^d))$. Each com- 13
14 ponent of \mathbf{h} is defined as 14

$$15 \quad h_j(x^j) = \begin{cases} h_{\bar{\delta}}(x^j) & \text{for } M_j = S^2, \\ \text{Log}_{\bar{x}^j}(x^j) & \text{otherwise,} \end{cases} \quad 16$$

17 where $\text{Log}_{\bar{x}^j}$ and $h_{\bar{\delta}}$ are defined in the Appendix and Section 4.1, respectively. 18

- 19 (3) Observe that $\mathbf{h}(x_1), \dots, \mathbf{h}(x_n) \in \mathbb{R}^{d_0}$ always have their mean at the origin. 19
20 Thus, the singular value decomposition of the $d_0 \times n$ data matrix $\mathbf{X} \equiv$ 20
21 $[\mathbf{h}(x_1) \cdots \mathbf{h}(x_n)]$ can be used for computation of the PCA. Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ 21
22 be the left singular vectors of \mathbf{X} corresponding to the largest m singular values. 22
23 (4) The k th principal arc is obtained by mapping the direction vectors \mathbf{v}_k onto M 23
24 by the inverse of h , which can be computed component-wise. 24

25 The principal arcs on M are not, in general, geodesics. Nor are they necessarily 25
26 circles, in the original marginal S^2 . This is because $h_{\bar{\delta}}$ and its inverse $h_{\bar{\delta}}^{-1}$ are 26
27 nonlinear transformations and, thus, a line on \mathbb{R}^2 may not be mapped to a circle 27
28 in S^2 . This is consistent with the fact that the principal components on a subset 28
29 of variables are different from projections of the principal components from the 29
30 whole variables. 30

31 Principal arc analysis for data on direct product manifolds often results in a 31
32 concise summary of the data. When we observe a significant variation along a 32
33 small circle of a marginal S^2 , that is most likely not a random artifact but, instead, 33
34 the result of a signal driving the circular variation. Nongeodesic variation of this 34
35 type is well captured by our method. 35
36

37 Principal arcs can be used to reduce the intrinsic dimensionality of M . Suppose 37
38 we want to reduce the dimension by k , where k can be chosen by inspection of the 38
39 scree plot. Then each data point x is projected to a k -dimensional submanifold M_0 39
40 of M in such a way that 40

$$41 \quad \mathbf{h}^{-1} \left(\sum_{i=1}^k \mathbf{v}_i \mathbf{v}_i^T \mathbf{h}(x) \right) \in M_0, \quad 41$$

42
43

1 where the \mathbf{v}_i 's are the principal direction vectors in \mathbb{R}^{d_0} , found by step 3 above. 1
 2 Moreover, the manifold M_0 can be parameterized by the k principal components 2
 3 z_1, \dots, z_k such that $M_0(z_1, \dots, z_k) = \mathbf{h}^{-1}(\sum_{i=1}^k z_i \mathbf{v}_i)$. 3
 4

5 4.1. *Choice of the transformation $h_{\bar{\delta}}$.* The transformation $h_{\bar{\delta}}: S^2 \rightarrow \mathbb{R}^2$ leads 5
 6 to an alternative representation of the data, which differs from the tangent space 6
 7 projection. The $h_{\bar{\delta}}$ transforms nongeodesic scatters along δ_1 to scatters along the x - 7
 8 axis, which makes a linear method like the PCA applicable. Among many choices 8
 9 of transformations that satisfy the three principles we stated, two methods are dis- 9
 10 cussed here. Recall that $\bar{\delta} = (\delta_1, \delta_2) = (\delta_1(\mathbf{c}, r), \mathbf{u})$. 10

11 *Projection.* The first approach is based on the projection of \mathbf{x} onto δ_1 , defined in 11
 12 (3), and a residual ξ . The signed distance from \mathbf{u} to $P_{\delta_1}x$, whose unsigned version 12
 13 is defined in (4), becomes the x -coordinate, while the residual ξ becomes the y - 13
 14 coordinate. This approach has the same spirit as the model for the circle class (5), 14
 15 since the direction of the signal is mapped to the x -axis, with the perpendicular 15
 16 axis for errors. 16

17 The projection $h_{\bar{\delta}}(\mathbf{x})$ that we define here is closely related to the spherical 17
 18 coordinate system. Assume $\mathbf{c} = \mathbf{e}_3$, and \mathbf{u} is at the Prime meridian (i.e., on the 18
 19 $x - z$ plane). For \mathbf{x} and its spherical coordinates (ϕ, θ) such that $\mathbf{x} = (x_1, x_2, x_3) =$ 19
 20 $(\cos \phi \sin \theta, \cos \phi \sin \theta, \cos \theta)$, 20

$$21 \quad (9) \quad h_{\bar{\delta}}(\mathbf{x}) = (\sin(r)\phi, \theta - \theta_{\mathbf{u}}), \quad 21$$

22 where $\theta_{\mathbf{u}} = \cos^{-1}(u_3)$ is the latitude of \mathbf{u} . The set of $h_{\bar{\delta}}(\mathbf{x}_i)$ has mean zero because 23
 24 the principal circle mean \mathbf{u} has been subtracted. 24

25 *Conformal map.* A conformal map is a function which preserves angles. We 25
 26 point out two conformal maps that can be combined to serve our purpose. See 26
 27 Chapter 9 of Churchill and Brown (1984) and Krantz (1999) for detailed discus- 27
 28 sions of conformal maps. A conformal map is usually defined in terms of complex 28
 29 numbers. Denote the extended complex plane $\mathbb{C} \cup \{\infty\}$ as \mathbb{C}^* . Let $\phi_{\mathbf{c}}: S^2 \rightarrow \mathbb{C}^*$ 29
 30 be the stereographic projection of the unit sphere when the point antipodal from 30
 31 \mathbf{c} is the projection point. Then $\phi_{\mathbf{c}}$ is a bijective conformal mapping defined for all 31
 32 S^2 that maps δ_1 as a circle centered at the origin in \mathbb{C}^* . The linear fractional trans- 32
 33 formation, sometimes called the Möbius transformation, is a rational function of 33
 34 complex numbers, that can be used to map a circle to a line in \mathbb{C}^* . In particular, 34
 35 we define a linear fractional transformation $f_{u^*}: \mathbb{C}^* \rightarrow \mathbb{C}^*$ as 35

$$36 \quad (10) \quad f_{u^*}(z) = \begin{cases} \frac{\alpha i(z - u^*)}{-z - u^*} & \text{if } z \neq -u^*, \\ \infty & \text{if } z = -u^*, \end{cases} \quad 36$$

37 where $u^* = \phi_{\mathbf{c}}(\mathbf{u})$, and α is a constant scalar. Then the image of δ_1 under $f_{u^*} \circ \phi_{\mathbf{c}}$ 40
 41 is the real axis, while the image of δ_2 is the imaginary axis. The mapping $h_{\bar{\delta}}: S^2 \rightarrow$ 41
 42 \mathbb{R}^2 is defined by $f_{u^*} \circ \phi_{\mathbf{c}}$ with the resulting complex numbers understood as mem- 42
 43 bers of \mathbb{R}^2 . Note that orthogonality of any two curves in S^2 is preserved by the $h_{\bar{\delta}}$ 43

PRINCIPAL ARC ANALYSIS

17




FIG. 7. Illustration of projection $h_{\bar{\delta}}$ [left, equation (9)] and conformal $h_{\bar{\delta}}$ [center, equation (10)] compared to a tangent plane projection at the geodesic mean (right) of the data in Figure 2c. The $h_{\bar{\delta}}$ maps the variation along δ_1 to the variation along the x-axis, while the tangent plane mapping fails to do so.

but the distances are not. Thus, we use the scale parameter α of the function f_{u^*} to match the resulting total variance of $h_{\bar{\delta}}(\mathbf{x}_i)$ to the geodesic variance of \mathbf{x}_i .

In many cases, both projection and conformal $h_{\bar{\delta}}$ give better representations than just using the tangent space. Figure 7 illustrates the image of $h_{\bar{\delta}}$ with the toy data set depicted in Figure 2c. The tangent space mapping is also plotted for comparison. The tangent space mapping leaves the curvy form of variation, while both $h_{\bar{\delta}}$'s capture the variation and lead to an elliptical distribution of the transformed data.

The choice between the projection and conformal mappings is a matter of philosophy. The image of the projection $h_{\bar{\delta}}$ is not all of \mathbb{R}^2 , while the image of the conformal $h_{\bar{\delta}}$ is all of \mathbb{R}^2 . However, in order to cover \mathbb{R}^2 completely, the conformal $h_{\bar{\delta}}$ can grossly distort the covariance structure of the data. In particular, the data points that are far from \mathbf{u} are sometimes overly diffused when the conformal $h_{\bar{\delta}}$ is used, as can be seen in the left tail of the conformal mapped image in Figure 7. The projection $h_{\bar{\delta}}$ does not suffer from this problem. Moreover, the interpretation of projection $h_{\bar{\delta}}$ is closely related to the circle class model. Therefore, we recommend the projection $h_{\bar{\delta}}$, which is used in the following data analysis.

5. Application to m-rep data. In this section an application of Principal Arc Analysis to the medial representation (m-rep) data example, introduced below in more detail, is described.

5.1. Medial representation. The m-rep gives an efficient way of representing 2- or 3-dimensional objects. The m-rep is constructed from the *medial axis*, which is a means of representing the middle of geometric objects. The medial axis of a 3-dimensional object is formed by the centers of all spheres that are interior to objects and tangent to the object boundary at two or more points. In addition, the




FIG. 8. (Left) An m -rep model with 3×5 grids of medial atoms. Each atom has its location (\mathbb{R}^3) and two equal-length spokes ($\mathbb{R}_+ \otimes S^2 \otimes S^2$). (Right) The implied surface of the m -rep model, showing a prostate shape.

medial description is defined by the centers of the inscribed spheres and by the associated vectors, called spokes, from the sphere center to the two respective tangent points on the object boundary. The medial axis is sampled over an approximately regular lattice and the elements of the lattice are called *medial atoms*. A medial atom consists of the location of the atom combined with two equal-length spokes, defined as a 4-tuple:

- location in \mathbb{R}^3 ;
- spoke direction 1, in S^2 ;
- spoke direction 2, in S^2 ;
- common spoke length in \mathbb{R}_+ ;

as shown in Figure 8. The size of the regular lattice is fixed for each object in practice. For example, the shape of a prostate is usually described by a 3×5 grid of medial atoms, across all samples. The collection of the medial atoms is called the medial representation (m -rep). An m -rep corresponds to a particular shape of prostate, and is a point in the m -rep space \mathcal{M} . The space of prostate m -reps is then $\mathcal{M} = (\mathbb{R}^3 \otimes \mathbb{R}_+ \otimes S^2 \otimes S^2)^{15}$, which is a 120-dimensional direct product manifold with 60 components. The m -rep model provides a useful framework for describing shape variability in intuitive terms. See Siddiqi and Pizer (2008) and Pizer et al. (2003) for detailed introduction to and discussion of this subject.

An important topic in medical imaging is developing segmentation methods of 3D objects from CT images; see Cootes and Taylor (2001) and Pizer et al. (2007). A popular approach is similar to a Bayesian estimation scheme, where the knowledge of anatomic geometries is used (as a prior) together with a measure of how the segmentation matches the image (as a likelihood). A prior probability distribution is modeled using m -reps as a means of measuring geometric atypicality of a segmented object. PCA-like methods (including PAA) can be used to reduce the dimensionality of such a model. A detailed description can be found in Pizer et al. (2007).

1 5.2. *Simulated m-rep object.* The data set partly plotted in Figure 1 is from the 1
 2 generator discussed in Jeong et al. (2008). It generates random samples of objects 2
 3 whose shape changes and motions are physically modeled (with some random- 3
 4 ness) by anatomical knowledge of the bladder, prostate and rectum in the male 4
 5 pelvis. Jeong et al. have proposed and used the generator to estimate the probabil- 5
 6 ity distribution model of shapes of human organs. 6

7 In the data set of 60 samples of prostate m-reps we studied, the major motion 7
 8 of prostate is a rotation. In some S^2 components, the variation corresponding to 8
 9 the rotation is along a small circle. Therefore, PAA should fit better for this type 9
 10 of data than principal geodesics. To make this advantage more clear, we also show 10
 11 results from a data set by removing the location and the spoke length information 11
 12 from the m-reps, the sample space of which is then $\{S^2\}^{30}$. 12

13 We have applied PAA as described in the previous section. The ratios μ/σ , 13
 14 estimated for the 30 S^2 components, are in general large (with minimum 21.2, 14
 15 median 44.1 and maximum 118), which suggests use of small circles to capture 15
 16 the variation. 16

17 Figure 9 shows the proportion of the cumulative variances, as a function of num- 17
 18 ber of components, from the Principal Geodesic Analysis (PGA) of Fletcher et al. 18
 19 (2004) and PAA. In both cases, the first principal arc leaves smaller residuals than 19
 20 the first principal geodesic. What is more important is illustrated in the scatter- 20
 21 plots of the data projected onto the first two principal components. The quadratic 21
 22 form of variation that requires two PGA components is captured by a single PAA 22
 23 component. 23

24 The probability distribution model estimated by principal geodesics is qualita- 24
 25 tively different from the distribution estimated by PAA. Although the difference in 25
 26 the proportion of variance captured is small, the resulting distribution from PAA 26
 27 is no longer elliptical. In this sense, PAA gives a convenient way to describe a 27
 28 nonelliptical distribution by, for example, a Normal density. 28
 29 29
 30 30

31 5.3. *Prostate m-reps from real patients.* We also have applied PAA to a 31
 32 prostate m-rep data set from real CT images. Our data consist of five patients' 32
 33 image sets, each of which is a series of CT scans containing prostate taken during 33
 34 a series of radiotherapy treatments [Merck (2008)]. The prostate in each image is 34
 35 manually segmented by experts and an m-rep model is fitted. The patients, coded 35
 36 as 3106, 3107, 3109, 3112 and 3115, have different numbers of CT scans (17, 12, 36
 37 18, 16 and 15, respectively). We have in total 78 m-reps. 37

38 The proportion of variation captured in the first principal arc is 40.89%, slightly 38
 39 higher than the 40.53% of the first principal geodesic. Also note that the estimated 39
 40 probability distribution model from PAA is different from that of PGA. In par- 40
 41 ticular, PAA gives a better separation of patients in the first two components, as 41
 42 depicted in the scatter plots (Figure 10). 42
 43 43




FIG. 9. (Top) The proportion of variances captured by the first few components of PAA are compared to those from PGA for the simulated prostate *m-reps*. (Bottom) Scatter plots of the data on $\{S^2\}^{30}$ show that the major variation is explained more concisely by the first principal arc.




FIG. 10. The scatter plots of the real prostate *m-rep* data. Different symbols represent different patients. PAA (right) gives a better separation of different patients in the first two components compared to PGA (left).

1 **6. Doubly iterative algorithm to find the least-squares small circle.** We 1
2 propose an algorithm to fit the least-squares small circle (1), which is a constrained 2
3 nonlinear minimization problem. This algorithm is best understood in two iterative 3
4 steps: The outer loop approximates the sphere by a tangent space; the inner loop 4
5 solves an optimization problem in the linear space, which is much easier than 5
6 solving (1) directly. In more detail, the $(k + 1)$ th iteration works as follows. The 6
7 sphere is approximated by a tangent plane at \mathbf{c}_k , the k th solution of the center of 7
8 the small circle. For the points on the tangent plane, any iterative algorithm to find 8
9 a least-squares circle can be applied as an inner loop. The solution of the inner 9
10 iteration is mapped back to the sphere and becomes the $(k + 1)$ th input of the outer 10
11 loop operation. One advantage of this algorithm lies in the reduced difficulty of 11
12 the optimization task. The inner loop problem is much simpler than (1) and the 12
13 outer loop is calculated by a closed-form equation, which leads to a stable and 13
14 fast algorithm. Another advantage can be obtained by using the *exponential map* 14
15 and *log map* (12) for the tangent projection, since they preserve the distance from 15
16 the point of tangency to the others, that is, $\rho(\mathbf{x}, \mathbf{c}) = \|\text{Log}_{\mathbf{c}}(\mathbf{x})\|$ for any $\mathbf{x} \in S^2$. 16
17 This is also true for radii of circles. The exponential map transforms a circle in 17
18 \mathbb{R}^2 centered at the origin with radius r to $\delta(\mathbf{c}, r)$. Thus, whenever (1) reaches its 18
19 minimum, the algorithm does not alter the solution. 19

20 We first illustrate necessary building blocks of the algorithm. A tangent plane 20
21 $T_{\mathbf{c}}$ at \mathbf{c} can be defined for any \mathbf{c} in S^2 , and an appropriate coordinate system of $T_{\mathbf{c}}$ 21
22 is obtained as follows. Basically, any two orthogonal complements of the direction 22
23 \mathbf{c} can be used as coordinates of $T_{\mathbf{c}}$. For example, when $\mathbf{c} = (0, 0, 1)' \equiv \mathbf{e}_3$, a coord- 23
24 inate system is given by \mathbf{e}_1 and \mathbf{e}_2 . For a general \mathbf{c} , let $q_{\mathbf{c}}$ be a rotation operator 24
25 on \mathbb{R}^3 that maps \mathbf{c} to \mathbf{e}_3 . Then a coordinate system for $T_{\mathbf{c}}$ is given by the inverse of 25
26 $q_{\mathbf{c}}$ applied to \mathbf{e}_1 and \mathbf{e}_2 , which is equivalent to applying $q_{\mathbf{c}}$ to each point of S^2 and 26
27 using $\mathbf{e}_1, \mathbf{e}_2$ as coordinates. 27

28 The rotation operator $q_{\mathbf{c}}$ can be represented by a rotation matrix. For $\mathbf{c} =$ 28
29 $(c_x, c_y, c_z)'$, the rotation $q_{\mathbf{c}}$ is equivalent to rotation through the angle $\theta =$ 29
30 $\cos^{-1}(c_z)$ about the axis $\mathbf{u} = (c_y, -c_x, 0)' / \sqrt{1 - c_z^2}$, whenever $\mathbf{c} \neq \pm \mathbf{e}_3$. When 30
31 $\mathbf{c} = \pm \mathbf{e}_3$, \mathbf{u} is set to be \mathbf{e}_1 . It is well known that a rotation matrix with axis 31
32 $\mathbf{u} = (u_x, u_y, u_z)'$ and angle θ in radians is, for $c = \cos(\theta)$, $s = \sin(\theta)$ and $v =$ 32
33 $1 - \cos(\theta)$, 33

$$(11) \quad R_{\mathbf{c}} = \begin{pmatrix} c + u_x^2 v & u_x u_y v - u_z s & u_x u_z v + u_y s \\ u_x u_y v + u_z s & c + u_y^2 v & u_y u_z v - u_x s \\ u_x u_z v - u_y s & u_y u_z v + u_x s & c + u_z^2 v \end{pmatrix},$$

38 so that $q_{\mathbf{c}}(\mathbf{x}) = R_{\mathbf{c}}\mathbf{x}$, for $\mathbf{x} \in \mathbb{R}^3$. 38

39 With the coordinate system for $T_{\mathbf{c}}$, we shall define the exponential map $\text{Exp}_{\mathbf{c}}$, 39
40 a mapping from $T_{\mathbf{c}}$ to S^2 , and the log map $\text{Log}_{\mathbf{c}} = \text{Exp}_{\mathbf{c}}^{-1}$. These are defined for 40
41 $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$ and $\mathbf{x} = (x_1, x_2, x_3)' \in S^2$, as 41

$$(12) \quad \text{Exp}_{\mathbf{c}}(\mathbf{v}) = q_{\mathbf{c}} \circ \text{Exp}_{\mathbf{e}_3}(\mathbf{v}), \quad \text{Log}_{\mathbf{c}}(\mathbf{x}) = \text{Log}_{\mathbf{e}_3} \circ q_{\mathbf{c}}(\mathbf{x}),$$

42

43

1 for $\theta = \cos^{-1}(x_3)$. See (18)–(19) in the [Appendix](#) for $\text{Exp}_{\mathbf{e}_3}$ and $\text{Log}_{\mathbf{e}_3}$. Note that
 2 $\text{Log}_{\mathbf{c}}(\mathbf{c}) = \mathbf{0}$ and $\text{Log}_{\mathbf{c}}$ is not defined for the antipodal point of \mathbf{c} .

3 Once we have approximated each \mathbf{x}_i by $\text{Log}_{\mathbf{c}}(\mathbf{x}_i) \equiv \tilde{\mathbf{x}}_i$, the inner loop finds the
 4 minimizer (\mathbf{v}, r) of

$$5 \quad (13) \quad \min \sum_{i=1}^n (\|\tilde{\mathbf{x}}_i - \mathbf{v}\| - r)^2, \quad 6$$

7
 8 which is to find the least-squares circle centered at \mathbf{v} with radius r . The general
 9 circle fitting problem is discussed in, for example, [Umbach and Jones \(2003\)](#) and
 10 [Chernov \(2010\)](#). This problem is much simpler than (1) because it is an uncon-
 11 strained problem and the number of parameters to optimize is decreased by 1.
 12 Moreover, optimal solution of r is easily found as

$$13 \quad (14) \quad \hat{r} = \frac{1}{n} \sum_{i=1}^n \|\tilde{\mathbf{x}}_i - \mathbf{v}\|, \quad 14$$

15 when \mathbf{v} is given. Note that for great circle fitting, we can simply put $\hat{r} = \pi/2$. Al-
 16 though the problem is still nonlinear, one can use any optimization method that
 17 solves nonlinear least squares problems. We use the Levenberg–Marquardt algo-
 18 rithm, modified by [Fletcher \(1971\)](#) [see Chapter 4 of [Scales \(1985\)](#) and Chapter 3
 19 of [Bates and Watts \(1988\)](#)], to minimize (13) with r replaced by \hat{r} . One can always
 20 use $\mathbf{v} = \mathbf{0}$ as an initial guess since $\mathbf{0} = \text{Log}_{\mathbf{c}}(\mathbf{c})$ is the solution from the previous
 21 (outer) iteration.

22 The algorithm is now summarized as follows:

- 23 (1) Given $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$, $\mathbf{c}_0 = \mathbf{x}_1$. 24
- 25 (2) Given \mathbf{c}_k , find a minimizer \mathbf{v} of (13) with r replaced by (14), with inputs
 26 $\tilde{\mathbf{x}}_i = \text{Log}_{\mathbf{c}_k}(\mathbf{x}_i)$. 26
- 27 (3) If $\|\mathbf{v}\| < \varepsilon$, then iteration stops with the solution $\hat{\mathbf{c}} = \mathbf{c}_k$, $r = \hat{r}$ as in (14). 27
 28 Otherwise, $\mathbf{c}_{k+1} = \text{Exp}_{\mathbf{c}_k}(\mathbf{v})$ and go to step 2. 28

29 Note that the radius of the fitted circle in $T_{\mathbf{c}}$ is the same as the radius of the
 30 resulting small circle. There could be many variations of this algorithm: as an
 31 instance, one can elaborate the initial value selection by using the eigenvector of
 32 the sample covariance matrix of \mathbf{x}_i 's, corresponding to the smallest eigenvalue as
 33 done in [Gray, Geiser and Geiser \(1980\)](#). Experience has shown that the proposed
 34 algorithm is stable and speedy enough. Gray, Geiser and Geiser proposed to solve
 35 (1) directly, which seems to be unstable in some cases. 35

36 The idea of the doubly iterative algorithm can be applied to other optimization
 37 problems on manifolds. For example, the geodesic mean is also a solution of a
 38 nonlinear minimization, where the nonlinearity comes from the use of the geodesic
 39 distance. This can be easily solved by an iterative approximation of the manifold to
 40 a linear space [See Chapter 4 of [Fletcher \(2004\)](#)], which is the same as the gradient
 41 descent algorithms [[Pennec \(1999\)](#), [Le \(2001\)](#)]. Note that the proposed algorithm,
 42 like other iterative algorithms, only finds one solution even if there are multiple
 43 solutions. 43

APPENDIX: SOME BACKGROUND ON DIRECT PRODUCT MANIFOLD

We give some necessary geometric background on direct product manifolds. Precise definitions and geometric discussions on a richer class of manifold, Riemannian manifold, can be found in Boothby (1986) and Helgason (2001).

A d -dimensional manifold can be thought of as a curved surface embedded in a Euclidean space of higher dimension d' ($\geq d$). The manifold is required to be smooth, that is, infinitely differentiable, so that a sufficiently small neighborhood of any point on the manifold can be well approximated by a linear space. The *tangent space* at a point p of a manifold M , $T_p M$, is defined as a linear space of dimension d which is tangent to M at p . The notion of distance on M is handled by a Riemannian metric, which is a metric of tangent spaces. In particular, the *geodesic distance function* $\rho_M(p, q)$ is roughly defined as the length of the shortest curve joining $p, q \in M$.

Now consider a direct product manifold $M = M_1 \otimes \dots \otimes M_m$. We restrict our attention to the case where each M_i is one of the simple manifolds S^1, S^2, \mathbb{R}_+ or \mathbb{R} , but most of the assertions below apply equally well to direct products of more general manifolds.

Geodesic distance function. The geodesic distance between $p \equiv (p^1, \dots, p^m)$ and $q \equiv (q^1, \dots, q^m)$ is defined by

$$\rho_M(p, q) = \left(\sum_{i=1}^m \rho_{M_i}^2(p^i, q^i) \right)^{1/2},$$

where each ρ_{M_i} is the geodesic distance function on M_i . The geodesic distance on S^1 is defined by the length of the shortest arc. Similarly, the geodesic distance on S^2 is defined by the length of the shortest great circle segment. The geodesic distance on \mathbb{R}_+ needs special treatment. In many practical applications, \mathbb{R}_+ represents a space of scale parameters. A desirable property for a metric on scale parameters is *scale invariance*, $\rho(rx, ry) = \rho(x, y)$ for any $x, y, r \in \mathbb{R}_+$. This can be achieved by differencing the logs, that is,

$$(15) \quad \rho_{\mathbb{R}_+}(x, y) = \left| \log \frac{x}{y} \right| \quad \text{for } x, y \in \mathbb{R}_+.$$

Finally, the geodesic distance on a simple manifold \mathbb{R} or \mathbb{R}^d is the Euclidean distance.

Geodesic mean and variance. The *geodesic mean* of a set of points in M , also referred to as the intrinsic mean, is also calculated component-wise. The *geodesic mean* of $x_1, \dots, x_n \in M$ is the minimizer in M of the sum of squared geodesic distances to the data. Thus, the geodesic mean is defined as

$$(16) \quad \bar{x} = \operatorname{argmin}_{x \in M} \frac{1}{n} \sum_{i=1}^n \rho_M^2(x, x_i).$$

1 In fact, each \bar{x}^i of $\bar{x} = (\bar{x}^1, \dots, \bar{x}^m)$ is the geodesic mean of $x_1^i, \dots, x_n^i \in M_i$. The
 2 geodesic mean of $\theta_1, \dots, \theta_n \in [0, 2\pi) \cong S^1$ is found by examining a candidate set
 3 consisting of

$$4 \quad (17) \quad \bar{\theta}_j = \frac{\sum_{i=1}^n \theta_i + 2j\pi}{n}, \quad j = 0, \dots, n-1,$$

5
 6 as in Moakher (2002). The geodesic mean in S^2 can be calculated by a full grid
 7 search or an iterative algorithm, as described in Section 6. The geodesic mean in
 8 \mathbb{R}_+ or \mathbb{R} is straightforward. Note that the geodesic mean may not be unique. How-
 9 ever, throughout the paper, we have assumed that the data have a unique geodesic
 10 mean which is true in most data analytic situations. Statistical investigation of the
 11 geodesic mean on manifolds can be found, for example, in Bhattacharya and Pa-
 12 trangenaru (2003, 2005) and Le and Kume (2000).

13 A related notion is *geodesic variance*. A sample geodesic variance is de-
 14 fined by the average squared geodesic distances to the geodesic mean, that is,
 15 $\frac{1}{n} \sum_{i=1}^n \rho_M^2(\bar{x}, x_i)$. When M is indeed the Euclidean space, the geodesic variance
 16 is the same as the total variance (the trace of the variance–covariance matrix).

17 *Mappings to tangent space.* The *exponential map* maps a point in $T_p M$ to M .
 18 The *log map* is the inverse exponential map whose domain is in M . For a direct
 19 product manifold M , the mappings are also defined component-wise. For S^1 , let
 20 $\theta \in \mathbb{R}$ denote an element of $T_p S^1$ where \mathbf{p} is set to be $(1, 0) \in S^1$ embedded in \mathbb{R}^2 .
 21 Then the exponential map is defined as

$$22 \quad \text{Exp}_{\mathbf{p}}(\theta) = (\cos \theta, \sin \theta).$$

23
 24 The corresponding log map of $\mathbf{x} = (x_1, x_2)$ is defined as $\text{Log}_{\mathbf{p}}(x) = \text{sign}(x_2) \cdot$
 25 $\arccos(x_1)$. For S^2 , let $\mathbf{v} = (v_1, v_2)$ denote a tangent vector in $T_p S^2$. Let $\mathbf{p} = \mathbf{e}_3$,
 26 then the exponential map $\text{Exp}_{\mathbf{p}} : T_p S^2 \rightarrow S^2$ is defined by

$$27 \quad (18) \quad \text{Exp}_{\mathbf{p}}(\mathbf{v}) = \left(\frac{v_1}{\|\mathbf{v}\|} \sin \|\mathbf{v}\|, \frac{v_2}{\|\mathbf{v}\|} \sin \|\mathbf{v}\|, \cos \|\mathbf{v}\| \right).$$

28
 29 This equation can be understood as a rotation of the base point \mathbf{p} to the direction
 30 of \mathbf{v} with angle $\|\mathbf{v}\|$. The corresponding log map for a point $\mathbf{x} = (x_1, x_2, x_3) \in S^2$
 31 is given by

$$32 \quad (19) \quad \text{Log}_{\mathbf{p}}(x) = \left(x_1 \frac{\theta}{\sin \theta}, x_2 \frac{\theta}{\sin \theta} \right),$$

33
 34 where $\theta = \arccos(x_3)$ is the geodesic distance from \mathbf{p} to x . Note that the antipodal
 35 point of p is not in the domain of the log map, that is, the domain of $\text{Log}_{\mathbf{p}}$ is
 36 $S^2/\{-\mathbf{p}\}$. In both S^1 and S^2 , \mathbf{p} can be set to be any point on S^1 or S^2 . The ex-
 37ponential and log map for an arbitrary T_p can be defined together with a rotation
 38operator, which is defined in Section 6 for the case of S^2 . The exponential map of
 39 \mathbb{R}_+ is defined by the standard real exponential function. The domain of the inverse
 40exponential map, the log map, is \mathbb{R}_+ itself. Finally, the exponential map on \mathbb{R}
 41is the identity map.
 42
 43

REFERENCES

- 1
2
3 BATES, D. M. and WATTS, D. G. (1988). *Nonlinear Regression Analysis and Its Applications*.
4 *Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics*. Wiley,
5 New York. [MR1060528](#)
- 6 BHATTACHARYA, R. and PATRANGENARU, V. (2003). Large sample theory of intrinsic and extrinsic
7 sample means on manifolds. I. *Ann. Statist.* **31** 1–29. [MR1962498](#)
- 8 BHATTACHARYA, R. and PATRANGENARU, V. (2005). Large sample theory of intrinsic and extrinsic
9 sample means on manifolds. II. *Ann. Statist.* **33** 1225–1259. [MR2195634](#)
- 10 BINGHAM, C. and MARDIA, K. V. (1978). A small circle distribution on the sphere. *Biometrika* **65**
11 379–389. [MR0513936](#)
- 12 BOOTHBY, W. M. (1986). *An Introduction to Differentiable Manifolds and Riemannian Geometry*,
13 2nd ed. Academic Press, Orlando, FL. [MR0861409](#)
- 14 CHERNOV, N. (2010). *Circular and Linear Regression: Fitting Circles and Lines by Least Squares*.
15 Chapman and Hall.
- 16 CHURCHILL, R. V. and BROWN, J. W. (1984). *Complex Variables and Applications*, 4th ed.
17 McGraw-Hill, New York. [MR0730937](#)
- 18 COOTES, T. F. and TAYLOR, C. J. (2001). Statistical models of appearance for medical image analy-
19 sis and computer vision. In *Proc. of SPIE Medical Imaging* **4322** 236–248.
- 20 COX, D. R. (1969). Some sampling problems in technology. In *New Developments in Survey Sam-*
21 *pling* (U. L. Johnson and H. Smith, eds.). Wiley, New York.
- 22 ELANDT, R. C. (1961). The folded normal distribution: Two methods of estimating parameters from
23 moments. *Technometrics* **3** 551–562. [MR0130738](#)
- 24 FISHER, N. I. (1993). *Statistical Analysis of Circular Data*. Cambridge Univ. Press, Cambridge.
25 [MR1251957](#)
- 26 FISHER, N. I., LEWIS, T. and EMBLETON, B. J. J. (1993). *Statistical Analysis of Spherical Data*.
27 Cambridge Univ. Press, Cambridge. [MR1247695](#)
- 28 FLETCHER, R. (1971). A modified Marquardt subroutine for non-linear least squares. Technical
29 Report No. AERE-R 6799.
- 30 FLETCHER, P. T. (2004). Statistical variability in nonlinear spaces: Application to shape analysis
31 and DT-MRI. PhD thesis, University of North Carolina at Chapel Hill.
- 32 FLETCHER, P. T., LU, C., PIZER, S. M. and JOSHI, S. (2004). Principal geodesic analysis for the
33 study of nonlinear statistics of shape. *IEEE Trans. Med. Imaging* **23** 995–1005.
- 34 GRAY, N. H., GEISER, P. A. and GEISER, J. R. (1980). On the least-squares fit of small and great
35 circles to spherically projected orientation data. *Math. Geol.* **12** 173–184. [MR0593408](#)
- 36 HASTIE, T. and STUETZLE, W. (1989). Principal curves. *J. Amer. Statist. Assoc.* **84** 502–516.
37 [MR1010339](#)
- 38 HELGASON, S. (2001). *Differential Geometry, Lie Groups, and Symmetric Spaces*. *Graduate Studies*
39 *in Mathematics* **34**. Amer. Math. Soc., Providence, RI. [MR1834454](#)
- 40 HUCKEMANN, S., HOTZ, T. and MUNK, A. (2010). Intrinsic shape analysis: Geodesic PCA for
41 Riemannian manifolds modulo isometric lie group actions. *Statist. Sinica* **20** 1–58. [MR2640651](#)
- 42 HUCKEMANN, S. and ZIEZOLD, H. (2006). Principal component analysis for Riemannian man-
43 ifolds, with an application to triangular shape spaces. *Adv. in Appl. Probab.* **38** 299–319.
[MR2264946](#)
- 44 JEONG, J.-Y., STOUGH, J. V., MARRON, J. S. and PIZER, S. M. (2008). Conditional-mean ini-
tialization using neighboring objects in deformable model segmentation. In *Proc. SPIE Medical*
Imaging **6914**.
- 45 JOHNSON, N. L. (1962). The folded normal distribution: Accuracy of estimation by maximum like-
lihood. *Technometrics* **4** 249–256. [MR0137221](#)
- 46 KARCHER, H. (1977). Riemannian center of mass and mollifier smoothing. *Comm. Pure Appl. Math.*
30 509–541. [MR0442975](#)

- 1 KRANTZ, S. G. (1999). *Handbook of Complex Variables*. Birkhäuser, Boston, MA. [MR1738432](#) 1
- 2 LE, H. (2001). Locating Fréchet means with application to shape spaces. *Adv. in Appl. Probab.* **33** 2
- 3 324–338. [MR1842295](#) 3
- 4 LE, H. and KUME, A. (2000). The Fréchet mean shape and the shape of the means. *Adv. in Appl.* 4
- 5 *Probab.* **32** 101–113. [MR1765168](#) 4
- 6 LEONE, F. C., NELSON, L. S. and NOTTINGHAM, R. B. (1961). The folded normal distribution. 5
- 7 *Technometrics* **3** 543–550. [MR0130737](#) 6
- 8 MARDIA, K. V. and GADSDEN, R. J. (1977). A circle of best fit for spherical data and areas of 7
- 9 vulcanism. *J. Roy. Statist. Soc. Ser. C* **26** 238–245. 8
- 10 MARDIA, K. V. and JUPP, P. E. (2000). *Directional Statistics. Wiley Series in Probability and Sta-* 9
- 11 *tistics*. Wiley, Chichester. [MR1828667](#) 9
- 12 MERCK, S. M., TRACTON, G., SABOO, R., LEVY, J., CHANEY, E., PIZER, S. M. and JOSHI, S. 10
- 13 (2008). Training models of anatomic shape variability. *Med. Phys.* **35** 3584–3596. 11
- 14 MOAKHER, M. (2002). Means and averaging in the group of rotations. *SIAM J. Matrix Anal. Appl.* 12
- 15 **24** 1–16 (electronic). [MR1920548](#) 13
- 16 PENNEC, X. (1999). Probabilities and statistics on Riemannian manifolds: Basic tools for geometric 14
- 17 measurements. In *International Workshop on Nonlinear Signal and Image Processing* **1** 194–198. 15
- 18 IEEE. 16
- 19 PIZER, S. M., FLETCHER, P. T., FRIDMAN, Y., FRITSCH, D. S., GASH, A. G., GLOTZER, J. M., 17
- 20 JOSHI, S., THALL, A., TRACTON, G., YUSHKEVICH, P. and CHANEY, E. L. (2003). De- 18
- 21 formable m-reps for 3D medical image segmentation. *Int. J. Comput. Vision* **55** 85–106. 19
- 22 PIZER, S. M., BROADHURST, R. E., LEVY, J., LIU, X., JEONG, J.-Y., STOUGH, J., TRACTON, G. 19
- 23 and CHANEY, E. L. (2007). Segmentation by posterior optimization of m-reps: Strategy and 20
- 24 results. Unpublished. 20
- 25 RIVEST, L.-P. (1999). Some linear model techniques for analyzing small-circle spherical data. 21
- 26 *Canad. J. Statist.* **27** 623–638. [MR1745827](#) 22
- 27 SCALES, L. E. (1985). *Introduction to Nonlinear Optimization*. Springer, New York. [MR0776609](#) 23
- 28 SIDDIQI, K. and PIZER, S. (2008). *Medial Representations: Mathematics, Algorithms and Applica-* 24
- 29 *tions*. Springer, New York. [MR2547467](#) 25
- 30 UMBACH, D. and JONES, K. N. (2003). A few methods for fitting circles to data. *IEEE Transactions* 26
- 31 *on Instrumentation and Measurement* **52** 1881–1885. 27
- 32 S. JUNG M. FOSKEY 28
- 33 J. S. MARRON DEPARTMENT OF RADIATION ONCOLOGY 29
- 34 DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA 30
- 35 AND OPERATIONS RESEARCH CHAPEL HILL, NORTH CAROLINA 27599 31
- 36 UNIVERSITY OF NORTH CAROLINA USA 32
- 37 CHAPEL HILL, NORTH CAROLINA 27599 E-MAIL: mark_foskey@unc.edu 33
- 38 USA 34
- 39 E-MAIL: sungkyu@email.unc.edu 35
- 40 marron@email.unc.edu 36
- 41 37
- 42 38
- 43 39
- 44 40
- 45 41
- 46 42
- 47 43

1 THE LIST OF SOURCE ENTRIES RETRIEVED FROM MATHSCINET 1

2 The list of entries below corresponds to the Reference section of your article and was retrieved
3 from MathSciNet applying an automated procedure. Please check the list and cross out those
4 entries which lead to mistaken sources. Please update your references entries with the data
5 from the corresponding sources, when applicable. More information can be found in the sup-
6 port page:

7 <http://www.e-publications.org/ims/support/mrhelp.html>. 7

8 BATES, D. M. AND WATTS, D. G. (1988). *Nonlinear regression analysis and its applications*.
9 Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics. John
10 Wiley & Sons Inc., New York. 1060528 (92f:62002) 10

11 BHATTACHARYA, R. AND PATRANGENARU, V. (2003). Large sample theory of intrinsic and ex-
12 trinsic sample means on manifolds. I. *Ann. Statist.* **31** 1–29. 1962498 (2004a:60069) 12

13 BHATTACHARYA, R. AND PATRANGENARU, V. (2005). Large sample theory of intrinsic and ex-
14 trinsic sample means on manifolds. II. *Ann. Statist.* **33** 1225–1259. 2195634 (2007j:60020) 14

15 BINGHAM, C. AND MARDIA, K. V. (1978). A small circle distribution on the sphere. *Biometrika* **65**
16 379–389. 513936 (80c:62064) 15

17 BOOTHBY, W. M. (1986). *An introduction to differentiable manifolds and Riemannian geometry*,
18 Second ed. Pure and Applied Mathematics, Vol. **120**. Academic Press Inc., Orlando, FL. 861409
19 (87k:58001) 18

19 Not Found! 19

20 CHURCHILL, R. V. AND BROWN, J. W. (1984). *Complex variables and applications*, Fourth ed.
21 McGraw-Hill Book Co., New York. 730937 (84k:30002) 20

21 Not Found! 21

22 Not Found! 22

23 ELANDT, R. C. (1961). The folded normal distribution: two methods of estimating parameters from
24 moments. *Technometrics* **3** 551–562. 0130738 (24 #A597b) 24

25 FISHER, N. I. (1993). *Statistical analysis of circular data*. Cambridge University Press, Cambridge.
26 1251957 (95b:62068) 25

27 FISHER, N. I., LEWIS, T., AND EMBLETON, B. J. J. (1993). *Statistical analysis of spherical*
28 *data*. Cambridge University Press, Cambridge. Revised reprint of the 1987 original. 1247695
29 (94h:62001) 28

29 Not Found! 29

30 Not Found! 30

31 Not Found! 31

32 GRAY, N. H., GEISER, P. A., AND GEISER, J. R. (1980). On the least-squares fit of small and great
33 circles to spherically projected orientation data. *J. Internat. Assoc. Math. Geol.* **12** 173–184.
34 593408 (82c:86007) 33

34 HASTIE, T. AND STUETZLE, W. (1989). Principal curves. *J. Amer. Statist. Assoc.* **84** 502–516.
35 1010339 (90i:62073) 35

36 HELGASON, S. (2001). *Differential geometry, Lie groups, and symmetric spaces*. Graduate Studies
37 in Mathematics, Vol. **34**. American Mathematical Society, Providence, RI. Corrected reprint of
38 the 1978 original. 1834454 (2002b:53081) 36

38 HUCKEMANN, S., HOTZ, T., AND MUNK, A. (2010). Intrinsic shape analysis: geodesic PCA for
39 Riemannian manifolds modulo isometric Lie group actions. *Statist. Sinica* **20** 1–58. 2640651 39

40 HUCKEMANN, S. AND ZIEZOLD, H. (2006). Principal component analysis for Riemannian mani-
41 folds, with an application to triangular shape spaces. *Adv. in Appl. Probab.* **38** 299–319. 2264946
42 (2007k:62102) 41

42 Not Found! 42

43 Not Found! 43

- 1 JOHNSON, N. L. (1962). The folded normal distribution: accuracy of estimation by maximum like- 1
2 lihood. *Technometrics* **4** 249–256. 0137221 (25 #677) 2
- 3 KARCHER, H. (1977). Riemannian center of mass and mollifier smoothing. *Comm. Pure Appl.* 3
4 *Math.* **30** 509–541. 0442975 (56 #1350) 4
- 5 KRANTZ, S. G. (1999). *Handbook of complex variables*. Birkhäuser Boston Inc., Boston, MA. 5
6 1738432 (2001a:30001) 5
- 7 LE, H. (2001). Locating Fréchet means with application to shape spaces. *Adv. in Appl. Probab.* **33** 6
8 324–338. 1842295 (2002d:60008) 7
- 9 LE, H. AND KUME, A. (2000). The Fréchet mean shape and the shape of the means. *Adv. in Appl.* 8
10 *Probab.* **32** 101–113. 1765168 (2001i:60015) 9
- 11 LEONE, F. C., NELSON, L. S., AND NOTTINGHAM, R. B. (1961). The folded normal distribution. 10
12 *Technometrics* **3** 543–550. 0130737 (24 #A597a) 11
- 13 Not Found! 11
- 14 MARDIA, K. V. AND JUPP, P. E. (2000). *Directional statistics*. Wiley Series in Probability and 12
15 Statistics. John Wiley & Sons Ltd., Chichester. Revised reprint of it Statistics of directional data 13
16 by Mardia [MR0336854 (49 #1627)]. 1828667 (2003b:62004) 14
- 17 Not Found! 14
- 18 MOAKHER, M. (2002). Means and averaging in the group of rotations. *SIAM J. Matrix Anal.* 15
19 *Appl.* **24** 1–16 (electronic). 1920548 (2004d:43008) 16
- 20 Not Found! 17
- 21 Not Found! 18
- 22 Not Found! 19
- 23 RIVEST, L.-P. (1999). Some linear model techniques for analyzing small-circle spherical data. 20
24 *Canad. J. Statist.* **27** 623–638. 1745827 21
- 25 SCALES, L. E. (1985). *Introduction to nonlinear optimization*. Springer-Verlag, New York. 776609 22
26 (86f:90136) 22
- 27 (2008). *Medial representations*. Computational Imaging and Vision, Vol. **37**. Springer, New York. 23
28 Mathematics, algorithms and applications, Edited by Kaleem Siddiqi and Stephen M. Pizer. 24
29 2547467 (2010g:65035) 25
- 30 Not Found! 26
- 31 27
- 32 28
- 33 29
- 34 30
- 35 31
- 36 32
- 37 33
- 38 34
- 39 35
- 40 36
- 41 37
- 42 38
- 43 39
- 44 40
- 45 41
- 46 42
- 47 43

1 META DATA IN THE PDF FILE 1

2 Following information will be included as pdf file Document Properties: 2

3
 4 **Title** : Principal Arc Analysis on direct product manifolds 4
 5 **Author** : Sungkyu Jung, Mark Foskey, J. S. Marron 5
 6 **Subject** : The Annals of Applied Statistics, 2010, Vol.0, No.00, 1- 6
 29
 7 **Keywords**: Principal Component Analysis, nonlinear dimension reduc- 7
 8 tion, manifold, folded Normal distribution, directional data, im- 8
 9 age analysis, medial representation 9

10
 11 THE LIST OF URI ADDRESSES 11

12
 13
 14 Listed below are all uri addresses found in your paper. The non-active uri addresses, if any, are 14
 15 indicated as ERROR. Please check and update the list where necessary. The e-mail addresses 15
 16 are not checked – they are listed just for your information. More information can be found in 16
 17 the support page: 17

18 <http://www.e-publications.org/ims/support/urihelp.html>. 18

19 200 <http://www.imstat.org/aoas/> [2:pp.1,1] OK 19
 20 200 <http://dx.doi.org/10.1214/10-AOAS370> [2:pp.1,1] OK 20
 21 200 <http://www.imstat.org> [2:pp.1,1] OK 21
 22 --- <mailto:sungkyu@email.unc.edu> [2:pp.23,23] Check skip 22
 23 --- <mailto:marron@email.unc.edu> [2:pp.23,23] Check skip 23
 24 --- mailto:mark_foskey@unc.edu [2:pp.23,23] Check skip 24
 25 200 <http://www.e-publications.org/ims/support/mrhelp.html> 25
 [2:pp.10000,10000] OK
 26 200 <http://www.e-publications.org/ims/support/urihelp.html> 26
 [2:pp.10002,10002] OK 27

28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43